

Physician Assistants: A Critical Role in Diagnosing and Treating Pesticide Poisonings:

Informing history taking and physical assessment of patients in ambulatory care settings by Advanced Practice Providers



Micah Bicker, PA-C



Rebecca Belloso, MPH

EPA Cooperative Agreement #X8-83935901 UC Davis in collaboration with Oregon State University

Objective 1 Clinicians will consid routine differential d

Clinicians will consider pesticide toxicity as part of routine differential diagnosis in specified populations

Objective 2

Clinicians will be made aware of resources for support of treatment of pesticide exposed patients

Objective 3

Clinicians will be made aware of resources for mandated reporting of pesticide toxicity where applicable

What are pesticides?

Any substance, or mixture of substances intended for preventing, repelling, destroying, or mitigating any pest. Examples:

> Fumigants

>Insecticides

> Herbicides

> Fungicides

➢ Rodenticides

➢ Antimicrobials



https://cfpub.epa.gov/giwiz/disinfectants/index.cfm



Why We Miss the Diagnosis

- We simply are not trained to recognize pesticide exposure and related symptoms.
- Exposure is often chronic and symptom onset may be insipid, impairing insight by patient/reporters that today's presentation/chronic problem may be related to toxic exposure
- Presenting symptoms may be attributed to other primary diagnosis (asthma, eczema, vertigo)
- Parents may be reluctant to share that a child/family member has been exposed

Barriers to Patient Reporting

- > Patients may simply be unaware of exposure
- Patients may not attribute presenting symptoms to exposures, especially if exposure is repeated/chronic
- Patients may not be wearing or have access to appropriate PPE
- Employees may be reluctant to report employers responsible for fear of retaliation/job loss
- Parents and caregivers may fear civil consequence for pediatric exposure, even if accidental
- Language/educational/cultural/communication barriers



Who experiences pesticide exposure?

- > Farm workers, in any agricultural setting
- > Factory/production workers (think meat processing/grain processing)
- Forestry workers
- > Hospitality employees, including restaurants
- Landscape/golf course employees
- > Anyone living in homes that may be insect/rodent infested
- Patients treated for lice/scabies/bedbugs
- Household members of any of the above



What will present clinically in the absence of an informed history?

- SKIN: contact dermatitis (look for patterning), burns, blisters, sores in/around the mouth, hyperhidrosis
- RESPIRATORY: asthmatic symptoms, vocal cord irritation including hoarseness and globus, salivation, rhinorrhea, lacrimation
- > GI: diarrhea, nausea/vomiting, anorexia, mucous in stool
- ≻GU: hematuria
- > NEUROLOGIC: headache, dizziness, lethargy, tremor, confusion, convulsions, seizures
- ENDOCRINE: amenorrhea, unexplained weight loss or weight gain, insulin resistance/increased blood sugars, symptoms of hypoglycemia without low blood sugar



When to take an environmental exposure history?

- If someone presents with respiratory s/s, or even mild hypersecretion, followed by ANY neurologic sequelae - even mild
- When a normally chronic illness presents as a new diagnosis. (New onset asthma in a 25 year old male without other atopy)
- When a known illness presents differently than prior (eczema presents in new distribution or is accompanied by desquamation)
- > When a contact dermatitis does not improve, or worsens with treatment
- With increased frequency of a patient presentation (a patient who normally is seen twice annually is in twice in a month)
- With endocrine disruption

What will stop me from thinking about pesticide toxicity? (When do I stop DDx?)

When a patient has an exacerbation of an underlying problem which is easily treated. For example – a patient presents with previous asthma, and an asthma exacerbation, gets oral steroids, and refills of SVN.



Long term effects of pesticide exposure: teaching points for patient education

- Inflammation and diseases related to inflammatory conditions such as vasculitis, and lupus.
- > Tumors of the prostate, pancreas, kidney and breast
- > Non-Hodgkin Lymphoma and Other Hematopoietic Cancers
- ➢ Parkinsonism
- > Endothelial inflammation and plaque deposition MI/CAD
- ≻ Obesity, DM2

Resources

www.pesticideresources.org/med



PESTICIDE TOXICITY: SIGNS AND SYMPTOMS

Pesticides are toxic by design. **Insecticides** and **rodenticides** are the pesticide types most commonly associated with acute pediatric poisoning. Warfarin type rodenticides in the form of pellets, grains or blocks are also a significant ingestion risk for young children. Many insecticide chemical classes designed to be neurotoxic for insect pests have been shown to be neurotoxic in humans, such as the cholinesterase inhibiting organophosphate and carbamate insecticides and the pyrethroids. Human toxicity varies by the pesticide product's active ingredient(s) and formulation (solvents, carriers). As such, signs and symptoms are broad and may include:



http://pesticideresources.org/med/resources/pubs.html

Pesticide Reporting Requirements



http://pesticideresources.org/med/reportingMap.html



Free Online CME

OEHHA's course for health care professionals on pesticiderelated illnesses and injuries.



education credit for RN and MD or a

general certificate.

http://pesticideresources.org/med/resources/training.html

Login

Recommended Clinical Reference

Recognition and Management of Pesticide Poisonings



http://pesticideresources.org//med/resources/rmpp.html



Case Study

LICE SHAMPOO OVEREXPOSURE

A 3 year old Hispanic male was brought to the local

AGE CONSIDERATION

- Vulnerability of Children
- Biological
- Behavioral
- Sociocultural Issues

POSSIBLE LEGAL ISSSUES

- Effectiveness of EO 13045
 EPA

 --Children's Health Policy
- --EPA BanNational Agenda

POSSIBLE SOCIAL ISSUES

- Substandard Housing
- Poor DietInadequate Health Care
- Exposure to Take Home Toxins

REPORTING ADVERSE EFFECTS

- Poison Control Center
- Other Community Resources
 Emergency Room
- Health Care Providers

emergency room by his Mother after experiencing a seizure in his home. Prior to this event, the child was discovered with the contents of a half-filled 2 ounce bottle of Lindane lice shampoo which he retrieved from the medicine cabinet in his home. His mother noted that the contents were on his cheek and chin, and he was trying to spit out the liquid; thus she induced vomiting by sticking her finger in his throat. She called her local poison control center and was told that the product can cause seizures; but that her child probably did not ingest enough to be at risk. She was advised to give him fluids to dilute the lindane that was ingested. About an hour later, the child

- fell on the bathroom floor and immediately began seizing
- (body stiffened, eyes rolled back in head, jaw shut tight,
- foamed at mouth); this seizure lasted for about 4 to 5

minutes.

This work is licensed under a <u>Creative Commons</u> Attribution-NonCommercial-ShareAlike 4.0 International License. © 2020 The Regents of the University of California,

• - - -

Davis campus. For information see this documer or contact PERC-MedSupport@ucdavis.edu. <u>Photo credit: "[mb] Medicine Cabinet"</u> by <u>Merrick Brown</u> is licensed under <u>CC BY-NC-SA 2.0</u> Original case study by:

PHEMED - PESTICIDE HEALTH EFFECTS MEDICAL EDUCATION DATABASE

ASSESSMENT AND PREVENTION

- Determining Exposure
- Acute Symptoms of Exposure
 Long Term Effects
- Long Term Ene
 Precautions

ABOUT LINDANE SHAMPOO

- Pharmacological Properties
- Purpose
- When to Use
- Contraindications

SOURCES OF

IngestionAbsorption

TREATMENT

- Neurological Problems
- Dermatological Problems
- Gastrointestinal Problems
- Long Term Effects

http://pesticideresources.org/med/resources/pubs.html

Future Directions

- Champion recruitment pilot
- Coding / billing sheet
- Conferences

Thank you!



micahbicker@hotmail.com

Rebecca A. Belloso, MPH

rabelloso@ucdavis.edu





Resources: Original PERC-med Works



Pesticide Illness & Family Medicine: How to Report in Your State

www.pesticideresources.org/med PERC-medSupport@ucdavis.edu

In the United States, pesticide usage is:

- ubiquitous
- estimated to total over 1.1 billion pounds annually (1).
- defined as any substance or mixture of substances intended for preventing, destroying, repelling or mitigating any pest.

Microorganisms, viruses and bacteria are included in the definition of pests.



High-risk occupations:

- both direct and indirect pesticide exposures
- the agriculture, hospitality, healthcare, building and grounds, environmental cleaning, forestry, and maintenance industries.

Concerns:

- Take home exposures,
- Pesticide residues may cling to a worker's skin, clothing, and work items creating potential for additional pesticide exposure for the worker's family.
- The COVID-19 pandemic has increased the use of antimicrobial pesticides (disinfectants) to help control the spread of SARS CoV-2. While antimicrobial pesticides are essential public health tools used in hospitals, schools, bathrooms and food preparation areas, there are risks related to their use.

Healthcare professionals play a critical role in recognizing and reporting pesticide-related illnesses. Pesticide illness surveillance information is used to inform policy and ideally prevent future pesticide illnesses. Additionally, it can identify atrisk populations, emerging pesticide issues, and occupations with increased exposure risks.



PERC-med developed an online interactive map and database covering the essentials of healthcare provider reporting of pesticide illnesses in all states and territories of the United States.

- Purpose: provide healthcare providers with an accurate and up to date tool to find information on their state's pesticide reporting requirements.
- Healthcare providers should be able to locate reporting requirements in their specific state(s) of practice.
- For states and territories that have mandatory or optional reporting, the tool describes who is required to report, what is defined as reportable, when the exposure needs to be reported by, and where it should be reported.

Reference

1. U.S. Environmental Protection Agency. Pesticides Industry Sales&Usage 2008–2012 Market Estimates, 2017.

The Pesticide Educational Resources Collaborative - Medical is a cooperative agreement (agreement #X-83935903) between the U.S. EPA's Office of Pesticide Programs and University of California Davis Extension, in collaboration with Oregon State University. PERC-med is funded by the U.S.EPA with the aim of helping medical professionals prevent, recognize and treat pesticide-related imeases and injuries.

Authors:

- Diana Simmes, MPH, Pesticide Medical Education Director, PERC-med Rebecca Belloso, MPH, Public Education Specialist, PERC-med
- · Kaci Buhl, MSc, Deputy Director, PERC-med
- Suzanne Forsyth MPH, Director, PERC-med
- Ann Marie O'Shea, Marketing Expert, PERC-med





What do Pesticides have to do with SARS-CoV-2 and COVID-19?

By: Diana R. Simmes, MPH and Rebecca Belloso, MPH PERC-med in Collaboration with the National Pesticide Information Center www.pesticideresources.org/med

Originally Published March 2020 in the Northwest Regional Primary Care Association (NWRPCA) Newsletter Updated November 2020

You may be surprised to learn that many infection control products commonly used in healthcare are actually pesticides.

More specifically, they are <u>antimicrobial</u> pesticides. Antimicrobial pesticides are essential public health tools because we use them in hospitals, schools, bathrooms and food preparation areas to prevent the spread of germs that can cause disease. So, what exactly constitutes a pesticide? A pesticide is defined as any substance or mixture of substances intended for preventing, destroying, repelling, or mitigating any pest. Microorganisms, bacteria and viruses are included in the definition of pests. Pesticides help protect our food, water, and health. However, there are always risks related to their use.